

# Progress towards simulating hyperspectral measurements to differentiate low-, middle-, and high-sensitivity models in CMIP5 (and other research highlights)

Daniel Feldman, John Paige, Xu Liu, William Collins,  
Yolanda Roberts, Peter Pilewskie

CLARREO 2014 Science Team Meeting

January 8, 2014

NASA Goddard Space Flight Center



# Presentation Outline

- Recap of OSSE tasks and accomplishments.
- MIROC5 vs HadGEM2-ES
  - Broadband
  - Hyperspectral
- Other research highlights
  - Arctic cloud feedbacks
  - Far infrared surface emissivity
- Future directions





# Proposed Tasks for the CLARREO SDT

- The Berkeley group has proposed to contribute the following to the CLARREO SDT:
  - Utilization of simulated CLARREO data to estimate change detection time in SW reflectance spectra
  - Interfacing different scenarios (varying forcings and feedbacks) of CCSM3 into the CLARREO OSSE framework.
  - Production of pan-spectral (SW+IR) OSSE spectra.
  - Production and analysis of spectra derived from different orbits (Roberts et al, AGU presentation; Jin et al, JGR, In Review).
  - Development and implementation of tools to produce OSSE spectra based on CMIP5 database.



# From CMIP5 OSSEs to CLARREO Science Questions




Taylor, 2009



# The Range of CMIP5 Climate Sensitivites


- Range of climate sensitivities in CMIP5 persists.
- How can comprehensive measurements narrow this range?
- Focus here is on MIROC5 vs HadGEM2-ES & RCP8.5 ( $2.72\text{--}4.59\text{ }^{\circ}\text{K}/2\times\text{CO}_2$ ).



Andrews, et al, 2012

# Formula for Change Detection

- Time-series differ where the secular trend in the signals exceeds the AR(1) noise envelope.
- Requires estimates of :
  - Natural variability.
  - Uncertainty in noise and trend estimation from a short time series.
  - Measurement uncertainty.



$$n^* \approx e^{\beta/\sqrt{M}} \left[ \frac{3.96\sigma_e}{|\omega_o|(1-\varphi)} \right]^{2/3} \left( 1 + \frac{\sigma_{meas}^2}{\sigma_v^2} \right)^{1/3}$$

Detection time

Scaling factor  
for length of  
record

Std dev of noise process

Measurement uncertainty

Feldman et al, 2013

Difference trend





AR(1) of noise process

Correlated natural variability







# All-Sky OLR Comparison

- Differential response in TWP OLR.
- Detectable with less than 10 years continuous data.



# Clear-Sky OLR Comparison





- Ts differences in Arctic lead to differences in OLRC.
- Detectable with less than 10 years of continuous data.









# All-Sky SW Comparison

- ITCZ and southern storm tracks.
- Detectable with  $\sim 20$  years of continuous data




# Clear-Sky SW Comparison

- Large differences in SW trends due to sea-ice response differences.
- Detectable with  $\sim 15$  years of continuous data.



# Porting the OSSE Framework to Sample CMIP5

Porting requires the addition of PCRTM and input homogenization



# Interfacing with CMOR-ized Data


- Thanks to the Climate Model Output Rewriter (CMOR), data in CMIP5 have a uniform format.
  - Clear-sky (condensate free)  $T_s$ ,  $T$ ,  $Q$ ,  $\text{CO}_2$ ,  $\text{O}_3$ ,  $\text{CH}_4$ ,  $\text{N}_2\text{O}$ .
  - Aerosols (Total aerosol, carbonaceous, dust, sea-salt, sulfate optical depths).
  - Clouds (Fractional area coverage, liq/ice distribution).
  - Surface boundary conditions (Snow cover, sea-ice cover, land fraction).
- These fields have been adapted externally to be consistent with the existing OSSE framework.





# Hyperspectral Results

- In Arctic, window bands in SW and LW differentiate models.
  - With individual channels, 10 year detection time without measurement uncertainty.
- In TWP, window bands + NIR H<sub>2</sub>O in SW and LW differentiate models.
  - With individual channels, 10 year detection time, without measurement uncertainty.
- Using PCs, we can exploit correlations of spectra and dipole responses to improve detection times.



Feldman, et al, GMD In Prep


# Intermodel Difference Detection

- A data record can differentiate 2 models in under 10 years based on vis window, nir H<sub>2</sub>O, and ir window.
  - Focus should be on few key locations because model response in these locations is so disparate.
  - Caveat: bootstrapping and validation with CFMIP quite challenging.
- For more subtle signals, how much longer are detection times?



# Diagnosis of Arctic Feedbacks in CMIP5



- Using APRP method of Taylor, diagnose cloud feedbacks in the Arctic from the CMIP5 database where albedo is stable despite loss of frozen surface extent.
- Determine how flux measurements can differentiate model outliers.
  - Requires TOA and surface fluxes.



Paige and Feldman, in Prep

# Other Research Highlights: $\epsilon_{\text{far-ir}}$


- For moist conditions, surface emissivity at wavelengths  $>15 \mu\text{m}$  does not impact radiation budget.
- For dry conditions,  $\epsilon_{\text{far-ir}}$  **DOES** impact radiation budget as atmosphere becomes more transparent through microwindows starting at  $550 \text{ cm}^{-1}$ .



Feldman, et al, PNAS Submitted

# $\epsilon_{\text{far-ir}}$ impacts OLR


- Below  $1 \text{ cm}^{-1}$  PWV, OLR is a strong function of  $\epsilon_{\text{far-ir}}$ .



Feldman, et al, PNAS Submitted

# Dry Conditions are Common



- AIRS indicates that high latitudes and altitudes are regularly dry enough such that  $\epsilon_{\text{far-ir}}$  matters.



Feldman, et al, PNAS Submitted

# Few Measurements of $\epsilon_{\text{far-ir}}$

- The published literature indicates that  $\epsilon_{\text{far-ir}}$  is a function of surface minerology.
- From U of Mich:  $\epsilon_{\text{far-ir}}$  is a function of snow grain size.
- Very few measurements, assumed to be 1.0 in climate models.
- Could be retrieved from single scenes with CLARREO far-IR spectrometer.
- No publication to justify extrapolation from mid-IR to far-IR (burden of proof is now on the extrapolator).




Cheng et al, 2013





# Climate Models Underspecify $\epsilon_{\text{far-ir}}$

- Climate models do not specify  $\epsilon_{\text{far-ir}}$  yet they are sensitive to it.
- Setting land  $\epsilon_{\text{far-ir}}$  to low value in Cheng et al, 2013 leads to:  
 $2^{\circ}\text{K}$   $\Delta T_s$ ,  $15 \text{ W/m}^2$   $\Delta \text{OLR}$ , and 15%  $\Delta \text{ice/snow fraction}$ , after only 15 years of integration.



Feldman, et al, PNAS Submitted



# Conclusions

- The OSSE is able to utilize CMOR-ized inputs to calculate SW reflectance and LW radiance
  - Validation activities are ongoing.
- Models of disparate climate sensitivities can be differentiated in under 10 years in TWP and Arctic.
- However, understanding interaction between clouds and frozen surfaces from conventional measurements takes many decades.
- CLARREO IR can produce far-IR emissivity immediately which cannot be extrapolated from mid-IR.



# Future Research Directions and Collaborations

- Radiometric validation of CMIP5 OSSE using CFMIP.
- Differentiation of low- and mid-sensitivity CMIP5 models.
- Pan-spectral PCA detection (with Yolanda).
- Detection analysis with AIRS spectral fluxes.
- Utilize spectral fingerprinting with OSSE data.

